

DIOXINS AND HEALTH

DIOXINS AND HEALTH

Including Other Persistent Organic Pollutants and Endocrine Disruptors

Third Edition

Edited by

ARNOLD SCHECTER

Department of Environmental and Occupational Health Sciences
University of Texas School of Public Health
Dallas, TX, USA

 **WILEY**

A JOHN WILEY & SONS, INC., PUBLICATION

Copyright © 2012 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permissions>.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Dioxins and health : including other persistent organic pollutants and endocrine disruptors / edited by Arnold Schecter. – 3rd ed.
p. ; cm.

Includes bibliographical references and index.

ISBN 978-0-470-60529-5 (cloth)

I. Schecter, Arnold.

[DNLM: 1. Dioxins—toxicity. 2. Environmental Pollutants—toxicity. WA 240]

LC-classification not assigned

615.9'512—dc23

2011033613

Printed in the United States of America.

ISBN: 9780470605295

10 9 8 7 6 5 4 3 2 1

To My Family

CONTENTS

PREFACE	xi
ACKNOWLEDGMENT	xvi
ABOUT THE EDITOR	xvii
CONTRIBUTORS	xix
LIST OF TABLES AND FIGURES	xxiii
1 PERSISTENT ORGANIC POLLUTANTS: AN OVERVIEW	1
<i>Daniele Wikoff, Lauren Fitzgerald, and Linda Birnbaum</i>	
2 TOXICITY EQUIVALENCE FACTORS FOR DIOXIN AND RELATED COMPOUNDS	37
<i>Michael DeVito</i>	
3 HISTORIC AND NEWER PERSISTENT ORGANIC POLLUTANTS IN FOOD	53
<i>Janice Huwe</i>	
4 FLAME RETARDANTS: POLYBROMINATED DIPHENYL ETHERS AND THEIR REPLACEMENTS	89
<i>Thomas F. Webster and Heather M. Stapleton</i>	
5 PHARMACOKINETICS OF 2,3,7,8-TETRACHLORODIBENZO-<i>p</i>-DIOXIN AND RELATED COMPOUNDS	109
<i>James R. Olson</i>	
	vii

6	IMMUNOMODULATION BY PERSISTENT ORGANIC POLLUTANTS	171
	<i>Robert W. Luebke, Jamie C. DeWitt, Dori R. Germolec, Keith D. Salazar, and Nancy I. Kerkvliet</i>	
7	DEVELOPMENTAL NEUROTOXICITY OF DIOXINS	193
	<i>G. Jean Harry and Pamela J. Lein</i>	
8	EPIDEMIOLOGICAL EVIDENCE ON THE HEALTH EFFECTS OF PERFLUOROOCCTANOIC ACID	229
	<i>Kyle Steenland, Tony Fletcher, and David A. Savitz</i>	
9	POLYBROMINATED DIBENZO-<i>p</i>-DIOXINS AND DIBENZOFURANS	255
	<i>Kurunthachalam Kannan, Chunyang Liao, and Hyo-Bang Moon</i>	
10	EPIDEMIOLOGICAL STUDIES ON CANCER AND EXPOSURE TO DIOXINS AND RELATED COMPOUNDS	303
	<i>Lennart Hardell and Mikael Eriksson</i>	
11	REPRODUCTIVE AND DEVELOPMENTAL EPIDEMIOLOGY OF DIOXINS	359
	<i>Anne Sweeney, Deborah J. del Junco, Marcella Warner, and Brenda Eskenazi</i>	
12	BISPHENOL A	381
	<i>Thaddeus T. Schug, Sarah A. Vogel, Laura N. Vandenberg, Joe M. Braun, Russ Hauser, Julia A. Taylor, Frederick S. vom Saal, and Jerold J. Heindel</i>	
13	PHTHALATES: HUMAN EXPOSURE AND RELATED HEALTH EFFECTS	415
	<i>John D. Meeker and Kelly K. Ferguson</i>	
14	THE SEVESO ACCIDENT	445
	<i>Angela Cecilia Pesatori and Pier Alberto Bertazzi</i>	
15	AGENT ORANGE: HEALTH AND ENVIRONMENTAL ISSUES IN VIETNAM, CAMBODIA, AND LAOS	469
	<i>Susan Hammond and Arnold Schecter</i>	
16	THE YUSHO AND YUCHENG RICE OIL POISONING INCIDENTS	521
	<i>Yoshito Masuda and Arnold Schecter</i>	

17 THE BINGHAMTON, NEW YORK, ELECTRICAL TRANSFORMER FIRE INCIDENT OF 1981	553
<i>Nancy Kim and Arnold Schechter</i>	
18 THE YUSHCHENKO DIOXIN POISONING: CHRONOLOGY AND PHARMACOKINETICS	567
<i>John Jake Ryan</i>	
19 POPs AND HUMAN HEALTH RISK ASSESSMENT	579
<i>David T. Szabo and Anne E. Loccisano</i>	
INDEX	619

PREFACE

This is the third edition of *Dioxins and Health*, which includes other persistent organic pollutants (POPs) and endocrine disruptors (EDs). The series began in 1994 with the publication of the first edition, written in an attempt to describe many complex issues related to health effects of dioxins and similar compounds in a less technical manner for those in different academic, scientific, and professional disciplines, as well as for the general public. In 2003, a second edition of *Dioxins and Health* was published to highlight scientific advances regarding dioxin exposure, health effects, and mechanisms of action. In organizing the second edition, we again found a number of world-class scientists to describe updated methods and findings from their disciplines. We decided to prepare a medium-sized volume with many references to introduce scientists and the general public to major issues related to dioxin rather than attempt to prepare an encyclopedic text.

With the development of what was originally meant to be a third edition of *Dioxins and Health*, we decided to include other POPs and endocrine disruptors, some similar to dioxins. We selected brominated flame retardants, such as polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecane (HBCD); perfluorinated chemicals (PFCs), such as perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS); and other endocrine disrupting chemicals similar to POPs in that they are found in human tissue at similar levels when measured, such as bisphenol A (BPA).

In the opening chapter, toxicologists Daniele Wikoff, Lauren Fitzgerald, and Linda Birnbaum introduce the readers to POPs, chemicals that are organic, toxic, persistent, and bioaccumulate. They review not only the classical POPs,

such as dioxins, polychlorinated biphenyls (PCBs), and dichlorodiphenyl trichloroethane (DTT), but also emerging POPs, such as PBDEs and HBCD. The authors also provide the reader with a generalized overview of the history of POPs, the concepts of biomagnification and biomonitoring, and provide an introduction to many of the chemicals discussed in later chapters.

Following this introduction, the concept of evaluating the toxicity of mixtures of dioxins and dioxin like compounds, including dibenzofurans and some PCBs, is presented by Michael DeVito, an NIH toxicologist, who also works with the World Health Organization (WHO) and other dioxin committees. This chapter introduces dioxin toxic equivalency factors (TEFs) and toxic equivalency (TEQ), an approach used to estimate the toxicity of dioxins and dioxin-like compounds. TEFs are derived by assigning an estimated order of magnitude toxicity value to each congener. These values are multiplied by the measured amount of each congener, then summed to produce one number as an estimate of total dioxin toxicity (TEQ). This approach has been a major step forward in dealing with chemical mixtures, such as dioxins and dibenzofurans, which act by a similar mechanism of action.

The issue of chemical contamination in food is addressed by Janice Huwe, a senior chemist with the United States Department of Agriculture (USDA), whose work on measuring dioxins, PBDEs, and other pollutants in U.S. food is well known. She reviews classical POPs, as well as emerging POPs to which humans may be exposed through consumption of food, including seafood, meat, poultry, dairy, eggs, and grains. She also notes differences in chemical residue levels found in foods occasionally due to local environmental contamination.

PBDEs and replacement flame retardants for PBDE mixtures are discussed by environmental scientist Thomas Webster and chemist Heather Stapleton. PBDEs, widely used flame retardants in the United States, are being phased out because of their toxicity; however, they are being replaced by newer flame retardants, many of which have unknown chemical composition and may also be toxic. Despite this phase out, PBDEs are found in human populations, biota, and the environment. Details of this ongoing problem and problems originating with the use of new flame retardants are described in this chapter.

James Olson, a toxicologist who has conducted research on dioxins and related compounds for decades, discusses the pharmacokinetics of selected POPs. Olson reviews studies that show the disposition and kinetics of POPs in both human and selected animal models, providing the reader with information regarding factors that can help explain biological conditions to these compounds, such as dose-dependent and time-dependent tissue distribution.

An overview of immunology and immunotoxicology precedes a discussion of specific immunological aspects of dioxins and other POPs in the chapter written by Robert Luebke, Jamie DeWitt, Dori Germolec, Keith Salazar, and Nancy Kerkvliet. Some effects on the immune system are believed to be responsible for the toxicity of these compounds. Findings are presented that show that the sensitivity of the immune system is high during the perinatal

period. Effects of POPs on the immune system include decreased ability to fight cancer and infectious diseases.

“Developmental Neurotoxicity of Dioxins,” by Jean Harry and Pamela Lein, introduces the reader to neurogenesis and provides an overview of *in vitro* and *in vivo* studies that show adverse effects on brain development. The authors focus on several possible aryl hydrocarbon receptor (AhR) dependent mechanisms of dioxin neurotoxicity and describe certain adverse neurologic health outcomes associated with prenatal PCB exposure, including decreased IQ, psychomotor skills, and deficits in visual recognition, and memory.

Kyle Steenland, Tony Fletcher, and David Savitz describe their studies on the health effects of PFOA and their epidemiological findings in exposed and comparison populations in the chapter titled “Epidemiologic Evidence on the Health Effects of Perfluorooctanoic Acid (PFOA).” In the past, PFCs have been used to repel stains or water in clothing and carpets, and have also been used in nonstick coatings found in some cooking utensils. Although there is considerable evidence of health effects in animal models, including several types of tumors, neonatal death loss, and toxic effects on the immune, hepatic, and endocrine systems, the effects of PFCs are not well characterized in humans. Production of some PFCs in the United States has ended, but because they are persistent, depot sources will remain in the environment for many years and will subsequently continue to contaminate members of the general population.

Chemists Kurunthachalam Kannan, Chunyang Liao, and Hyo-Bang Moon introduce polybrominated dioxins (PBDDs) and dibenzofurans (PBDFs). These compounds are not as well known as their chlorinated cousins (PCDDs and PCDFs) because of the complex and costly analytical procedures needed to study them, and also because they are frequently found at lower levels than PCDD/Fs. However, they are believed to cause similar health effects as the chlorinated dioxins and dibenzofurans. The authors provide background information on these chemicals and their origins. They also discuss similarities and differences between PBDD/Fs and PCDD/Fs.

Cancer has long been a concern from exposure to dioxins, PCBs, and related POPs. Lennart Hardell and Mikael Eriksson review the literature on this subject, including their own groundbreaking research on dioxins and cancers in humans. The authors review epidemiological studies showing associations between POPs exposure and cancers, including soft tissue sarcomas, malignant lymphomas, multiple myeloma, leukemia, nasal and nasopharyngeal cancers, hepatic cancer, and gastric cancer. The cancers considered by many to have the strongest associations with dioxin exposure are non-Hodgkin’s lymphoma and soft tissue sarcoma, first linked to dioxin exposure in studies by these authors.

Reproductive and developmental effects of dioxins and other POPs are discussed by epidemiologists Anne Sweeney, Deborah del Junco, Marcella Warner, and Brenda Eskenazi. Inability to conceive, spontaneous abortion, altered menstrual periods, and endometriosis are possible reproductive effects

of POP exposure. The authors review many studies regarding reproductive outcomes, including the Seveso Women's Health Study, paternal dioxin exposure, and environmental dioxin exposure.

The chapter on BPA, by Thaddeus Schug, Sarah Vogel, Laura Vandenberg, Joe Braun, Russ Hauser, Julia Taylor, Frederick vom Saal, and Jerold Heindel, provides an introduction to this endocrine disrupter. It has been commonly found in can linings and plastic bottles, such as baby bottles and water bottles. Although BPA has a short half-life in humans and is not persistent, it is sometimes considered similar to POPs because of its frequent intake. At the time of this book's preparation, there is a debate concerning what levels of BPA intake should be considered safe. Some scientists feel present government levels are too permissive. Epidemiological studies are not consistent with respect to BPA health; however, animal studies have shown obesity, decreased semen quality, cardiovascular disease, endocrine disruption, and cancer as health effects following BPA exposure.

Phthalate health findings are presented by John D. Meeker and Kelly K. Ferguson. Phthalates are found in many products, including perfumes, deodorants, cosmetics, and other personal care products, to which most in industrialized countries are regularly exposed. Phthalates are ubiquitous and have proven to be a challenge when measured in tissue, food, and other media. Laboratory contamination with phthalates has made analysis a major challenge. Data exists on phthalate levels in biota but quantification remains difficult. The endocrine-disrupting characteristics of phthalates may lead to reproductive and developmental effects in men and women, obesity, cancers, and possibly altered sex differentiation from fetal exposure.

The next chapters of the text focus on a number of incidents characterized by the presence of certain POPs. Epidemiologists Angela Cecilia Pesatori and Pier Alberto Bertazzi describe the well-known 1976 dioxin incident in Seveso, Italy. An explosion at a chemical factory released 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on a residential population. In contrast to Vietnam and most other dioxin contaminated locations, the site of the Seveso contamination has been converted into a public park with precautions to reduce or prevent dioxin release from the soil underlying the park. A number of epidemiological studies, assessing cancer and other health effects, have been performed from this incident.

Susan Hammond, a social activist, and Arnold Schechter, a public health physician-researcher, each long active in Vietnam work, discuss Agent Orange and its dioxin contaminant, 2,3,7,8-TCDD, sprayed in Vietnam between 1962 and 1971. They describe major issues, including persistence of dioxin in the environment, people, and wildlife; social and legal issues arising from this controversial spraying; remediation efforts; and monetary issues involved in remediation.

Two of the most definitive incidents documenting human health damage from dioxin-like chemicals have been those of the rice oil contamination

incidents in Japan and Taiwan, known as Yusho and Yucheng, respectively. Certain batches of rice oil used for cooking were contaminated with PCBs, PCDFs, and other toxicants in the 1960s in Japan and 1970s in Taiwan. Approximately 2000 people were classified as victims by government agencies for each incident. Professor Yoshito Masuda, the chemist who discovered and characterized PCDFs in the Japanese incident, describes both incidents in his Yusho and Yucheng chapter, which is coauthored by physician and scientific collaborator Arnold Schecter.

A fifth event involving dioxins is the 1981 Binghamton (New York) State Office Building (BSOB) incident, in which an electrical panel caught fire and subsequently heated PCBs and chlorinated benzenes in a nearby basement electrical transformer. The heated soot moved up air ducts and stairways to contaminate the entire office building. It took 13 years and approximately \$50,000,000 to sufficiently clean the building and allow reoccupancy. This incident contributed substantially to the concept, validation, and implementation of TEFs and TEQs. In addition, the incident stimulated major advances in detection of individual dioxin and dibenzofuran congeners in environmental samples, human adipose tissue (fat tissue), and blood at low levels and hence, biomonitoring. The study of this incident unexpectedly found that all persons sampled, including those from the general population, had dioxins and dibenzofurans in their bodies. Before that time, it was believed that only certain individuals had dioxin contamination. Toxicologist Nancy Kim and physician Arnold Schecter, then working for the State of New York Department of Health and for the local county (Broome County) health department, respectively, review this incident from their unique perspectives as responders beginning at the time of the incident.

John Jake Ryan, a Canadian chemist formerly with Health Canada, describes the last major incident discussed in this book: The dioxin poisoning of former Ukrainian president Viktor Yushchenko, one of the better-known dioxin incidents worldwide. Victor Yushchenko was a politician running for the presidency of Ukraine when he mysteriously became ill. This was found to be a deliberate poisoning with pure 2,3,7,8-TCDD, the most toxic dioxin. Prior to the presidential election, he attended a late-night dinner with political opponents and abruptly became ill with symptoms now known to be from dioxin poisoning. Mr. Yushchenko continued his presidential campaign and became president of Ukraine despite health problems related to the dioxin poisoning.

The final chapter in the book, written by toxicologists David Szabo and Anne Loccisano, provides the reader with a summary of human health risk assessment focusing on many of the POPs that are currently of substantial concern. It serves to bring together in one self-contained chapter an overview of the chemicals covered in previous chapters, and some new chemicals, from the perspective of how to deal with and assess human health risk from these toxic compounds.

This book presents information on historical, social, and scientific aspects of dioxins and other related POPs. It endeavors to provide the reader with a concise yet relatively comprehensive overview of major compounds currently of interest with sufficient references to help the reader pursue issues of special interest in greater depth.

Dallas, Texas
2012

ARNOLD SCHECTER

ACKNOWLEDGMENT

This book was prepared with the expert assistance of Noor Malik, Tyra Gent, James Miller, and Sarah Smith.

ABOUT THE EDITOR



Arnold Schecter is a public health and preventive medicine physician who specializes in research in biomonitoring to estimate exposure to various toxic chemicals. Collaborations include scientists in the United States, Germany, Finland, Canada, Vietnam, Cambodia, Laos, Japan, Russia, Israel, and other countries.

He began his work with dioxins by documenting dioxin and dibenzofuran formation from an electrical transformer fire in Binghamton, New York,

collaborating with various expert chemists when they demonstrated that patterns found in fat tissue and blood of exposed workers were similar to those found in environmental samples from the fire. The U.S. Environmental Protection Agency (EPA) later decided to ban the use of PCBs in electrical transformers in public buildings based on this and other incidents.

Later, he studied dioxins from Agent Orange in Vietnamese and Americans. His research in Vietnam began in 1984 and continues to the present. His research documented elevated dioxin levels from Agent Orange in Vietnam, including elevated dioxin levels in Vietnamese food, blood, and milk and environmental samples from defoliation agents used in previous decades. The studies also documented dioxins from other sources in Vietnamese human tissues. He and his colleagues documented elevated dioxin levels in workers as much as 30–40 years after exposure.

More recently, with other scientists and physicians, his group documented PBDE levels in the milk of nursing American women. They also noted the levels were the highest to date by orders of magnitude compared with other then-published reports from various countries. This and other works were used to reduce PBDE manufacture and use. His recent work includes the study of human and food levels of HBCD, BPA, phthalates, and other persistent organic pollutants (POPs) and endocrine disruptors (EDs). Studies of home-based electrical waste recycling workers in less developed countries are some of the newest projects.

He enjoys communicating with a broad audience through various media. He served on EPA and ATSDR on dioxin and related chemicals expert panels and also served as a (dioxin) Special Expert at the National Institute of Environmental Health Sciences (NIEHS/NIH). He has served on several scientific editorial boards. His training was at the University of Chicago, Howard Medical School, and Columbia University School of Public Health. His employment has been on the faculties of several State University of New York campuses and New Jersey Medical School, as well as with several community-based health programs. He currently is a professor at the University of Texas School of Public Health, Dallas Campus.

CONTRIBUTORS

PIER ALBERTO BERTAZZI, Department of Occupational and Environmental Health, Università degli Studi di Milano and Fondazione IRCCS Ca' Granda—Ospedale Maggiore Policlinico, Milan, Italy

LINDA BIRNBAUM, National Cancer Institute/National Institute of Environmental Health Sciences, Durham, NC

JOE M. BRAUN, Department of Environmental Health and Department of Epidemiology, Harvard School of Public Health, Boston, MA

DEBORAH J. DEL JUNCO, Centers for Translational Injury Research and Clinical and Translational Sciences, University of Texas Health Science Center, Houston, TX

MICHAEL DEVITO, National Toxicology Program, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, NC

JAMIE C. DEWITT, Department of Pharmacology and Toxicology, Brody School of Medicine, East Carolina University, Greenville, NC

MIKAEL ERIKSSON, Department of Oncology, Skane University Hospital and Lund University, Lund, Sweden

BRENDA ESKENAZI, Center for Environmental Research and Children's Health, School of Public Health, University of California, Berkeley, CA

KELLY K. FERGUSON, Department of Environmental Sciences, University of Michigan School of Public Health, Ann Arbor, MI

LAUREN FITZGERALD, ToxStrategies, Austin, TX

TONY FLETCHER, London School of Hygiene and Tropical Medicine, University of London, London, UK

DORI R. GERMOLEC, Toxicology Branch, National Toxicology Program, National Institute of Environmental Health Sciences, Research Triangle Park, NC

SUSAN HAMMOND, War Legacies Project, Chester, VT

LENNART HARDELL, Department of Oncology, University Hospital, Örebro, Sweden

G. JEAN HARRY, Laboratory of Toxicology and Pharmacology, National Institute of Environmental Health Sciences, Research Triangle Park, NC

RUSS HAUSER, Department of Environmental Health and Department of Epidemiology, Harvard School of Public Health, Boston, MA

JEROLD J. HEINDEL, Division of Extramural Research and Training, Cellular, Organ and Systems Pathobiology Branch, National Institute of Environmental Health Sciences, Research Triangle Park, NC

JANICE HUWE, Agricultural Research Service, United States Department of Agriculture, Fargo, ND

KURUNTHACHALAM KANNAN, Wadsworth Center, New York State Department of Health, and Department of Environmental Health Sciences, School of Public Health, State University of New York at Albany, Albany, NY

NANCY I. KERKVLiet, Department of Environmental & Molecular Toxicology, Oregon State University, Corvallis, OR

NANCY KIM, Valatie, NY

PAMELA J. LEIN, Department of Molecular Biosciences, School of Veterinary Medicine, University of California, Davis, Davis, CA

CHUNYANG LIAO, Wadsworth Center, New York State Department of Health, and Department of Environmental Health Sciences, School of Public Health, State University of New York at Albany, Albany, NY

ANNE E. LOCCISANO, The Hamner Institutes for Health Sciences, Research Triangle Park, NC

ROBERT W. LUEBKE, Cardiopulmonary and Immunotoxicology Branch, U.S. Environmental Protection Agency, Research Triangle Park, NC

YOSHITO MASUDA, Daiichi College of Pharmaceutical Sciences, Fukuoka, Japan

JOHN D. MEEKER, Department of Environmental Sciences, University of Michigan School of Public Health, Ann Arbor, MI

HYO-BANG MOON, Department of Environmental Marine Sciences, College of Science and Technology, Hanyang University, Ansan, Republic of Korea

JAMES R. OLSON, University at Buffalo, State University of New York (SUNY), Buffalo, New York

ANGELA CECILIA PESATORI, Department of Occupational and Environmental Health, Università degli Studi di Milano and Fondazione IRCCS Ca' Granda—Ospedale Maggiore Policlinico, Milan, Italy

JOHN JAKE RYAN, Health Canada, Health Products and Food Branch, Bureau Chemical Safety, Ottawa, Ontario, Canada

KEITH D. SALAZAR, Cardiopulmonary and Immunotoxicology Branch, U.S. Environmental Protection Agency, Research Triangle Park, NC

DAVID A. SAVITZ, Department of Community and Preventive Medicine, Mount Sinai School of Medicine, New York

ARNOLD SCHECTER, Department of Environmental and Occupational Health Sciences, University of Texas School of Public Health, Dallas Regional Campus, Dallas, TX

THADDEUS T. SCHUG, Cellular, Organ and Systems Pathobiology Branch, Division of Extramural Research and Training, National Institute of Environmental Health Sciences, Research Triangle Park, NC

HEATHER M. STAPLETON, Nicholas School of Environment and Earth Sciences, Duke University, Durham, NC

KYLE STEENLAND, Rollins School of Public Health, Emory University, Atlanta, GA

ANNE SWEENEY, Texas A&M Health Science Center School of Rural Public Health, Texas A&M University, College Station, TX

DAVID T. SZABO, Curriculum in Toxicology, University of North Carolina, Chapel Hill, NC

JULIA A. TAYLOR, Division of Biological Sciences, University of Missouri, Columbia, MO

LAURA N. VANDENBERG, Center for Regenerative and Developmental Biology, Department of Biology, Tufts University, Medford, MA

SARAH A. VOGEL, Johnson Family Foundation, New York, NY

FREDERICK S. VOM SAAL, Division of Biological Sciences, University of Missouri, Columbia, MO

MARCELLA WARNER, Center for Environmental Research and Children's Health, School of Public Health, University of California, Berkeley, Berkeley, CA

THOMAS F. WEBSTER, School of Public Health, Boston University, Boston, MA

DANIELE WIKOFF, ToxStrategies, Austin, TX

LIST OF TABLES AND FIGURES

CHAPTER 1 PERSISTENT ORGANIC POLLUTANTS: AN OVERVIEW

Table 1.1.	Chemicals Currently Controlled under the Stockholm Convention	3
Table 1.2.	Measurements of Selected POPs in the General U.S. Population as Reported by NHANES	5
Figure 1.1.	Chemical structures of selected organochlorine pesticides.	6
Figure 1.2.	General chemical structures for PCDDs (left) and PCDFs (right).	7
Figure 1.3.	Chemical structure of polychlorinated biphenyls (PCBs).	7
Figure 1.4.	TCDD's hypothesized modes of action in site-specific carcinogenesis in rodents. See text for details. In each instance, the solid arrows depict pathways that are well established and are associated with low uncertainty. The dashed arrows represent connections that are less established and are associated with higher uncertainty. Adapted from reference [12].	9
Figure 1.5.	Chemical structure of polybrominated biphenyls (PBBs).	12
Figure 1.6.	Chemical structure of polybrominated diphenyl ethers (PBDEs).	14

- Figure 1.7. Chemical structures of HBCD stereoisomers. 19
 Figure 1.8. Chemical structure of PFOA (left) and PFOS (right). 20

CHAPTER 2 TOXICITY EQUIVALENCE FACTORS FOR DIOXIN AND RELATED COMPOUNDS

- Table 2.1. Toxic Equivalency Factors (TEFs) 43

CHAPTER 3 HISTORIC AND NEWER PERSISTENT ORGANIC POLLUTANTS IN FOOD

- Table 3.1. Mean Levels and Ranges of Selected POPs in Fish and Seafood on a Wet Weight (ww) Basis 56
 Table 3.2. Mean Levels and Ranges of Selected POPs in Meat and Poultry Samples 63
 Table 3.3. Mean Levels and Ranges of Selected POPs in Eggs and Dairy Products 69
 Table 3.4. Mean Levels and Ranges of Selected POPs in Cereals, Fruit, Vegetables, and Miscellaneous Foods 75

CHAPTER 4 FLAME RETARDANTS: POLYBROMINATED DIPHENYL ETHERS AND THEIR REPLACEMENTS

- Table 4.1. Timeline for PBDEs and Other Flame Retardants 92
 Figure 4.1. Chemical structures for PBDEs and several alternate flame retardants now detected in humans and/or the environment. 90
 Figure 4.2. Application of the source to disease paradigm to PBDEs. 94

CHAPTER 5 PHARMACOKINETICS OF 2,3,7,8-TETRACHLORODIBENZO-*p*-DIOXIN AND RELATED COMPOUNDS

- Table 5.1. Net Gastrointestinal Absorption or Excretion of PCDDs, PCDFs, and PCBs in Humans and Dependence on Congener-Specific Blood Lipid Levels 113
 Table 5.2. 2,3,7,8-Substituted PCDDs and PCDFs in Human Liver and Adipose Tissue 121
 Table 5.3. Elimination Constants and Half-Lives of Various 2,3,7,8-Substituted PCDDs and PCDFs in Hepatic and Adipose Tissue of Marmoset Monkeys 123
 Table 5.4. Half-Life Estimates for 2,3,7,8-TCDD and Related Compounds in Humans 131
 Table 5.5. Reference Half-Life Values and Model Parameters in Humans for PCDDs and PCDFs (Adapted from Milbrath et al.) 143

Table 5.6.	Reference Half-Life Values and Model Parameters in Humans for Dioxin-Like PCBs (Adapted from Milbrath et al.)	146
Table 5.7.	Impact of Various Parameters on Estimates of Daily Intake of 2,3,7,8-TCDD Utilizing Equation 5.7	151

CHAPTER 8 EPIDEMIOLOGICAL EVIDENCE ON THE HEALTH EFFECTS OF PERFLUOROCTANOIC ACID

Table 8.1.	Changes in Cholesterol in Relation to Changes in PFOA Levels	234
------------	--	-----

CHAPTER 9 POLYBROMINATED DIBENZO-*p*-DIOXINS AND DIBENZOFURANS

Table 9.1.	Predicted Values of Log Octanol–Water Partition Coefficients (K_{OW}), Octanol–Air Partition Coefficients (K_{OA}) and Air–Water Partition Coefficients (K_{AW}) of PBDDs and PBDFs	257
Table 9.2.	Selected Laboratory Thermolysis Studies Reporting Concentrations of PBDD/Fs from Brominated Flame Retardant Standards	260
Table 9.3.	Summary of PBDD/F Concentrations Reported in Various Environmental Compartments, Biota, and Humans	271
Table 9.4.	Relative Potency Values of Chlorinated, Brominated, and Mixed Substituted Dioxins and Furans (PXCD/Fs) Relative to 2,3,7,8-TCDD Using DR-CALUX Assay	284
Figure 9.1.	Generalized structures of PCDD/Fs and PBDD/Fs.	256
Figure 9.2.	Thermolysis of decabromodiphenyl ether (deca-BDE).	261
Figure 9.3.	Condensation of naturally occurring bromophenols to brominated dibenzo- <i>p</i> -dioxins (adapted from Haglund).	267
Figure 9.4.	Average homologue composition (%) of PBDD/Fs in (a) water ($n = 15$; Litten et al.), (b) air ($n = 63$ for Taiwan; Wang et al., $n = 5$ for Japan; Hayakawa et al., and $n = 25$ for China, Li et al.), (c) dust ($n = 2$ for house dust; Tue et al., $n = 5$ for TV cabinets; Takigami et al., $n = 10$ for e-waste facility in Vietnam; Tue et al., and $n = 5$ for e-waste facility in China; Ma et al.), (d) soil ($n = 10$, Ma et al.), (e) marine sediment ($n = 6$ for Hong Kong and $n = 8$ for South Korea; Terauchi et al.) and (f) biota ($n = 9$ for fish, $n = 3$ for crab, and $n = 6$ for bivalve; Miyake et al.) reported from previous studies. The error bars denote the standard deviation for individual PBDD/F homologues. D4, tetra-BDDs; D5, penta-BDDs; D6, hexa-BDDs; D7, hepta-BDDs; D8, octa-BDD; F4, tetra-BDFs; F5, penta-BDFs; F6, hexa-BDFs; F7, hepta-BDFs; F8, octa-BDF.	280

CHAPTER 10 EPIDEMIOLOGICAL STUDIES ON CANCER AND EXPOSURE TO DIOXINS AND RELATED COMPOUNDS

Table 10.1.	Odds Ratios and 95% Confidence Intervals (in Parentheses) in Swedish Case–Control Studies on Soft-Tissue Sarcoma	306
Table 10.2.	Mantel–Haenszel Odds Ratios and 90% Confidence Intervals Adjusted by Study for STS among Persons Exposed to All Dioxins, TCDD, and Dioxins Other than TCDD in Four Case–Control Studies Involving 434 Cases and 948 Controls	307
Table 10.3.	Odds Ratios and 95% Confidence Intervals (in Parentheses) in Swedish Case–Control Studies on Malignant Lymphoproliferative Diseases	312
Table 10.4.	Number of White Male NHL Cases and Controls and Odds Ratios by Days Per Year of Exposure to 2,4-D in Kansas and Nebraska	314

CHAPTER 12 BISPHENOL A

Table 12.1.	BPA Interactions with Various Nuclear Receptors	384
Figure 12.1.	Chemical structures of BPA, DES, and estradiol.	383

CHAPTER 13 PHTHALATES: HUMAN EXPOSURE AND RELATED HEALTH EFFECTS

Table 13.1.	Phthalates and Urinary Metabolites (Adapted from CDC)	418
Table 13.2.	Health Outcomes in Infants and Children Associated with Phthalate Exposure (Adapted from Swan, Meeker, and Bornehag and Nanberg)	421
Table 13.3.	Health Outcomes in Adults Associated with Phthalate Exposure (Adapted from Swan, Meeker et al., and Jaakola and Knight)	430

CHAPTER 14 THE SEVESO ACCIDENT

Table 14.1.	TCDD Soil Measurements, Serum TCDD, and Serum TEQ Levels in Samples of Subjects Resident in the Seveso Area	450
-------------	---	-----

Table 14.2.	Summary of the Endpoints Examined in the Seveso Women Health Study	453
Table 14.3.	Neonatal b-TSH Levels in Children Born between 1994 and 2005 to Women from Zone A, B, and the Reference Area	455
Table 14.4.	Mortality, 1976–2001, from Selected Cancer Causes	457
Table 14.5.	Cancer Incidence, 1977–1996, in the Seveso Population, Aged 20–74 Years, Resident in the Area at the Accident Time	458
Figure 14.1.	Main steps in exposure-response relationship assessment.	447
Figure 14.2.	Map of the Seveso area.	448
Figure 14.3.	Infant sex ratios and exposure status of fathers stratified by fathers' ages at the accident time.	452
Figure 14.4.	Parkwood opened in July 1996.	461

CHAPTER 15 AGENT ORANGE: HEALTH AND ENVIRONMENTAL ISSUES IN VIETNAM, CAMBODIA, AND LAOS

Table 15.1.	Use of Herbicides in Vietnam 1961–1971 in Liters Used	472
Table 15.2.	Number of Hectares Sprayed with Herbicides	472
Table 15.3.	Estimates of TCDD Contamination in the 2,4,5-T	473
Figure 15.1.	Map of the herbicide spray missions between 1965 and 1971. Each gray line represents one of the aerial spray missions, and includes all types of tactical herbicides used in Operation Ranch Hand (Agents Purple, Pink, Green, Orange, Blue, and White). <i>Source:</i> U.S. Department of the Army and Hatfield Consultants.	475
Figure 15.2.	Some of the areas of Vietnam that have been demonstrated to have elevated levels of dioxin in soil or sediment. Da Nang, Phu Cat, and Bien Hoa Airbase require mitigation measures to reduce TEQ in the soil and sediment to the Vietnamese government's standard of 1000ppt TEQ for soils and 150ppt TEQ for sediment. A Luoi (A So Special Forces Base) was just under 1000ppt TEQ for soil; however, the community living directly on the hotspot was relocated. Limited testing has been done on the other areas identified on the map, most of which were former Ranch Hand bases or herbicide storage sites. These TCDD levels were found at a given time, but do not represent all levels measured at these sites. Levels as high as over 1,000,000ppt TCDD in soil was detected at Bien Hoa.	486

CHAPTER 16 THE YUSHO AND YUCHENG RICE OIL POISONING INCIDENTS

Table 16.1.	Concentrations of PCBs, PCDFs, and PCQs in Yusho Oil and Kanechlor-400	524
Table 16.2.	Concentrations of PCDD, PCDF, and PCB Congeners and TEQs in the Rice Oil, Calculated by the WHO 1998 and 2005	525
Table 16.3.	Mean Estimated Intakes of Rice Oil and TEQs by Yusho Patients (Range in Parentheses)	526
Table 16.4.	Concentrations of PCB Congeners in the Blood of Yusho Patients, Sampled in 1983	527
Table 16.5.	Concentrations of HO-PCBs and PCBs in Yusho Patients ($n = 9$)	529
Table 16.6.	Concentrations of PCDF Congeners in the Tissues and Blood of Yusho Patients	530
Table 16.7.	Concentrations (ppb in Blood) and Half-Life (Years) of PCBs and PCDFs in the Blood of Yucheng and Yusho Patients ($n = 3$ and $n = 5$) After the Onset	533
Table 16.8.	Percent Distribution of Symptoms of Yusho Patients Examined Before October 31, 1968	535
Table 16.9.	Types and Concentrations of Blood PCBs and Incidence of Clinical Symptoms among 72 Yusho Patients from April 1973 to March 1974	536
Figure 16.1.	Gas chromatograms of PCB fractions on the 5% SE-30 glass column (3 mm \times 2 m): (a) Kanechlor-400; (b) Kanemi rice oil; (c,d) blood of typical Yusho patients; (e) blood from a type C Yusho patient; and (f) blood from a normal control. Adapted from Reference 16.	523
Figure 16.2.	Yusho intake, regulation, and personal intakes of TEQ. Adapted from Reference 60.	542
Figure 16.3.	Concentration trend of PCBs and TEQ in Yusho and control in Japan, 1968–2007. Adapted from Reference 15.	543
Figure 16.4.	Normal levels of hormones and concentrations of PCBs, PCDFs, and DDE in the blood of Yusho patients (severe: black; common: gray; controls: white). Adapted from Matsuda.	544

CHAPTER 17 THE BINGHAMTON, NEW YORK, ELECTRICAL TRANSFORMER FIRE INCIDENT OF 1981

Table 17.1.	Calculated 2,3,7,8-TCDD-Equivalent Concentrations of the Binghamton Soot for Various Dose-Related End Points Following Subchronic Exposure to Guinea Pigs	558
-------------	---	-----

**CHAPTER 18 THE YUSHCHENKO DIOXIN POISONING:
CHRONOLOGY AND PHARMACOKINETICS**

Table 18.1.	Chronology of the Yushchenko Incident	570
Table 18.2.	Symptoms, Signs, and Treatment of Yushchenko Poisoning	571
Table 18.3.	Pharmacokinetic Findings Over 3 Years on Yushchenko by the Swiss-Ukraine Group	573
Table 18.4.	Pharmacokinetics of the Two Highest Known Exposed Humans to 2,3,7,8-TCDD: Yushchenko and Vienna Patient No. 1	574
Figure 18.1.	Photographs of Victor Yushchenko before poisoning (A), 3 months (B), and 3.5 years after poisoning with 2,3,7,8-tetrachlorodibenzo- <i>p</i> -dioxin; taken from article published by Sorg et al. in <i>Lancet</i> 374(9696)1179–1185 (2009), fig. 2, p. 1180. Reproduced with permission of Elsevier Ltd., Oxford, UK.	568

CHAPTER 19 POPs AND HUMAN HEALTH RISK ASSESSMENT

Table 19.1.	Original and Emerging POPs as Defined by the Stockholm Convention	580
Table 19.2.	Risk Values Derived by Various Agencies for HCHs	585
Table 19.3.	Risk Values Derived by Various Agencies for Chlordecone	591
Table 19.4.	Risk Values Derived by Various Agencies for HBB	593
Table 19.5.	Risk Values Derived by Various Agencies for PeCB	594
Table 19.6.	Risk Values Derived by Various Agencies for PFOS	596
Table 19.7.	Risk Values Derived by Various Agencies for PBDEs	598
Table 19.8.	Risk Values Derived by Various Agencies for HBCD	601